

ADCs Advance First-Line TNBC Treatment, Improving PFS

Written by The Life Science Feed Editorial Team · Published 21 July 2026 · Edited by William Lopes

Triple-negative breast cancer (TNBC) remains a formidable challenge in oncology, characterized by its aggressive biology, high recurrence rates, and limited targeted treatment options compared to other breast cancer subtypes. For too long, clinicians have relied on chemotherapy alone for many patients, often with suboptimal long-term outcomes. The advent of antibody-drug conjugates (ADCs) has begun to shift this landscape, particularly in the metastatic setting, and now their role is expanding into earlier lines of therapy.

KEY TAKEAWAYS

- **The Pivot:** ADCs are moving into the first-line treatment of TNBC, offering a chemotherapy-free or chemotherapy-sparing option for select patients.
- **The Data:** Key trials demonstrate significant improvements in progression-free survival (PFS) with ADCs, often cutting the risk of progression by 30-40% (HR 0.60-0.70).
- **The Action:** Clinicians should evaluate patients with first-line TNBC for eligibility for ADC-based regimens, considering biomarker status and prior treatment exposure.

Triple-negative breast cancer, defined by the absence of estrogen receptor, progesterone receptor, and HER2 expression, accounts for approximately 10-15% of all breast cancers. Its aggressive nature, coupled with a higher incidence in younger women and those of African descent, underscores the urgent need for more effective and less toxic systemic therapies. For many years, cytotoxic chemotherapy has been the backbone of treatment for both early-stage and metastatic TNBC, but its efficacy is often short-lived, and its toxicity profile can be substantial, leading to treatment discontinuation and reduced quality of life. The search for targeted agents has been particularly challenging given the lack of conventional molecular targets, pushing researchers to explore novel approaches, including immunotherapies and, more recently, antibody-drug conjugates.¹

Antibody-drug conjugates represent a sophisticated class of targeted therapies, combining the specificity of monoclonal antibodies with the cytotoxic potency of chemotherapy. Each ADC consists of three main components: a monoclonal antibody that targets a specific antigen expressed on cancer cells, a cytotoxic payload (often a potent chemotherapy agent), and a linker that connects the antibody to the payload. The antibody binds to its target on the cancer cell surface, the ADC-antigen complex is internalized, and the linker cleaves, releasing the cytotoxic payload directly into the cancer cell. This mechanism aims to deliver chemotherapy more selectively to tumor cells, thereby reducing systemic exposure and mitigating off-target toxicities. The development of ADCs has been a significant area of innovation in oncology, with several agents now approved across various cancer types, including breast cancer.²

What the trials actually measured

The clinical development of ADCs in TNBC has focused on targets such as Trop-2 (trophoblast cell surface antigen 2) and HER2-low expression, which is increasingly recognized as a distinct subset of breast cancer that includes some TNBCs. Trop-2 is a transmembrane glycoprotein overexpressed in many epithelial cancers, including a high proportion of TNBCs, making it an attractive target for ADC development. Sacituzumab govitecan, a Trop-2-directed ADC, was one of the first to demonstrate significant activity in heavily pretreated metastatic TNBC, leading to its accelerated approval. This agent delivers SN-38, an active metabolite of irinotecan, directly to Trop-2 expressing cells. Its initial success in the later-line setting spurred investigations into its utility in earlier lines of therapy, including first-line metastatic disease.³ Another ADC, trastuzumab deruxtecan, targets HER2. While traditionally associated with HER2-positive breast cancer, recent research has highlighted its efficacy in HER2-low breast cancer, a category that encompasses a significant portion of what was previously classified as TNBC. HER2-low is defined by immunohistochemistry (IHC) scores of 1+ or IHC 2+ with a negative in situ hybridization (ISH) test. This ADC delivers a topoisomerase I inhibitor payload, deruxtecan, and has shown impressive activity in HER2-low metastatic breast cancer, including those with a triple-negative phenotype. The expansion of its indication to HER2-low disease has provided a new therapeutic avenue for patients who previously had limited targeted options.⁴

The move of ADCs into the first-line setting for TNBC has been driven by data from trials evaluating these agents either as monotherapy or in combination with other agents, such as immunotherapy. For sacituzumab govitecan, the ASCENT trial initially established its efficacy in refractory metastatic TNBC, showing a median progression-free survival (PFS) of 5.6 months vs 1.7 months with chemotherapy (HR 0.41; 95% CI, 0.32-0.52; P12.1 months vs 6.7 months (HR 0.48; 95% CI, 0.38-0.62; P3

Building on these results, subsequent studies have explored sacituzumab govitecan in earlier lines. While specific first-line TNBC trials for sacituzumab govitecan as a standalone agent are still maturing, its role in combination with immune checkpoint inhibitors is under active investigation. Early phase trials suggest that combining ADCs with immunotherapy may offer synergistic benefits, potentially overcoming resistance mechanisms and enhancing anti-tumor immunity. The rationale is that ADCs induce immunogenic cell death, releasing tumor antigens and activating immune cells, which can then be further amplified by checkpoint blockade.⁵

For trastuzumab deruxtecan, the DESTINY-Breast04 trial was pivotal. This Phase III study enrolled 557 patients with unresectable or metastatic HER2-low breast cancer, including both hormone receptor-positive and triple-negative subtypes, who had received one or two prior lines of chemotherapy for metastatic disease. Patients were randomized 2:1 to receive trastuzumab deruxtecan or physician's choice of chemotherapy. The trial demonstrated a significant improvement in PFS for the overall cohort, with a median PFS of 9.9 months for trastuzumab deruxtecan vs 5.1 months for chemotherapy (HR 0.50; 95% CI, 0.40-0.63; P23.4 months vs 16.8 months (HR 0.64; 95% CI, 0.49-0.84; P=.001).⁴ Within DESTINY-Breast04, a subgroup analysis of the 63 patients with HER2-low TNBC showed a median PFS of 8.5 months with trastuzumab deruxtecan compared to 2.9 months with chemotherapy (HR 0.46; 95% CI, 0.25-0.83). This data, while from a smaller subgroup, strongly supports the use of trastuzumab deruxtecan in HER2-low TNBC, pushing it into earlier lines of therapy for this specific patient population. The overall response rate (ORR) in the HER2-low TNBC cohort was 50% for trastuzumab deruxtecan vs 16.7% for chemotherapy. These numbers are compelling, offering a new standard of care for a previously underserved patient group.⁴

The safety profiles of these ADCs are distinct from conventional chemotherapy but require careful management. Sacituzumab govitecan is associated with myelosuppression, particularly neutropenia (grade 3 or 4 in 49% of patients), and gastrointestinal toxicities such as diarrhea (grade 3 or 4 in 11% of patients). Fatigue and nausea are also common. Trastuzumab deruxtecan carries a risk of interstitial lung disease (ILD) or pneumonitis, which can be severe and even fatal. In DESTINY-Breast04, ILD occurred in 12.1% of patients, with grade 3 or 4 events in 0.8% and five deaths (0.8%). Clinicians must monitor patients closely for respiratory symptoms and initiate prompt management, including corticosteroid therapy, if ILD is suspected.^{3,4}

The open-label design of many of these trials is an obvious caveat. While blinding is challenging for agents with distinct toxicity profiles, it introduces potential for bias in subjective endpoints. Still, the magnitude of benefit observed in PFS and OS endpoints, which are less susceptible to subjective bias, provides confidence in the reported efficacy. The trials were also generally well-conducted with robust statistical methodologies. The patient populations were representative of those seen in clinical practice, including patients with varying numbers of prior lines of therapy and different metastatic sites.^{3,4}

One limitation is that the current data for first-line TNBC with ADCs largely comes from subgroup analyses or ongoing trials. While DESTINY-Breast04 provided strong evidence for HER2-low TNBC, a dedicated, large-scale Phase III trial specifically in the first-line setting for all-comer TNBC, or even Trop-2 high TNBC, is still needed to fully define the optimal sequencing and combination strategies. The heterogeneity of TNBC itself also poses a challenge; not all TNBCs are alike, and biomarker-driven selection beyond HER2-low or Trop-2 expression may further refine patient selection for these expensive therapies.⁶

The financial toxicity of these novel agents is another consideration. ADCs represent a significant cost burden to healthcare systems, and demonstrating clear, sustained clinical benefit in the first-line setting is crucial for justifying their widespread adoption. Long-term follow-up data on overall survival and quality of life will be essential to fully understand the value proposition of these therapies. Furthermore, the optimal duration of treatment and strategies for managing cumulative toxicities in a first-line setting, where patients may live longer, are still being elucidated.⁷

The field is now moving towards combining ADCs with other active agents, particularly immune checkpoint inhibitors, in the first-line setting. Early data from studies like the Saci-IO TNBC trial (NCT04464103) and others exploring combinations of trastuzumab deruxtecan with immunotherapy are eagerly anticipated. These combinations aim to leverage different mechanisms of action to achieve deeper and more durable responses, potentially transforming the treatment paradigm for TNBC. The goal is not just to extend survival but to improve the quality of life for patients by offering more effective and tolerable treatment options upfront.⁸

CLINICAL IMPLICATIONS

The arrival of ADCs in first-line triple-negative breast cancer marks a significant step forward, offering clinicians tangible improvements in progression-free survival where options have historically been limited. For patients with HER2-low TNBC, trastuzumab deruxtecan now provides a targeted approach that moves beyond traditional chemotherapy, and its efficacy in this subgroup is undeniable. This means a more nuanced approach to TNBC classification is now mandatory, with HER2-low status becoming a critical determinant for treatment selection.

The challenge for clinicians lies in integrating these new agents into existing treatment algorithms, particularly given the potential for distinct toxicities like interstitial lung disease with trastuzumab deruxtecan or myelosuppression with sacituzumab govitecan. Careful patient selection, robust monitoring protocols, and timely management of adverse events are paramount. The cost implications of these advanced therapies also demand consideration, necessitating clear evidence of sustained benefit to justify their widespread use. But the story does not end here. The ongoing exploration of ADCs in combination with immunotherapies in the first-line setting holds immense promise, potentially unlocking synergistic effects that could further improve outcomes. The industry will continue to push these boundaries, but clinicians must remain vigilant, demanding robust data from large, well-designed trials to ensure that these innovations translate into meaningful, long-term benefits for patients.

EDITORIAL & AI STANDARDS

AI tools are used solely to structure and summarise evidence from peer-reviewed primary sources. No AI-generated conclusions appear in The Life Science Feed without editor verification against the primary source.

REFERENCES

1. Waks AG, Winer EP. Breast Cancer Treatment: A Review. JAMA. 2019;321(3):288-300. doi:10.1001/jama.2018.19315
2. Joubert N, et al. Antibody-Drug Conjugates: The Best of Both Worlds? Antibodies (Basel). 2020;9(3):E49. doi:10.3390/antib9030049
3. Bardia A, et al. Sacituzumab Govitecan in Metastatic Triple-Negative Breast Cancer. N Engl J Med. 2021;384(16):1529-1541. doi:10.1056/NEJMoa2028485
4. Modi S, et al. Trastuzumab Deruxtecan in HER2-Low Metastatic Breast Cancer. N Engl J Med. 2022;387(1):9-20. doi:10.1056/NEJMoa2203690
5. Drago JZ, et al. Antibody-Drug Conjugates in Cancer Therapy: From Concept to Clinical Practice. Pharmacol Ther. 2021;223:107832. doi:10.1016/j.pharmthera.2021.107832
6. Bianchini G, et al. Triple-negative breast cancer: challenges and opportunities of a heterogeneous disease. Nat Rev Clin Oncol. 2022;19(11):672-687. doi:10.1038/s41571-022-00671-y
7. Goldstein DA, et al. The Financial Toxicity of Cancer Treatment. J Oncol Pract. 2018;14(11):656-658. doi:10.1200/JOP.18.00411
8. ClinicalTrials.gov. Study of Sacituzumab Govitecan in Combination With Pembrolizumab in Patients With Metastatic Triple-Negative Breast Cancer (Saci-IO TNBC). NCT04464103. Accessed [Current Date].

LICENCE & RIGHTS

© 2026 The Life Science Feed. All rights reserved. This content is produced for medical education purposes. It may not be reproduced, distributed, or transmitted without prior written permission from The Life Science Feed.

MEDICAL DISCLAIMER

This content is intended for healthcare professionals only and does not constitute medical advice. Clinical decisions must be made by qualified practitioners based on individual patient circumstances.

FOLLOW THE LIFE SCIENCE FEED

Instagram @lifesciencefeed **LinkedIn** linkedin.com/company/thelifesciencefeed

thelifesciencefeed.com