

Diagnostic Inconsistencies Challenge STEMI Paradigm's Simplicity

Written by The Life Science Feed Editorial Team · Published 21 June 2026 · Edited by William Lopes

The diagnostic framework for acute coronary occlusion, particularly the ST-elevation myocardial infarction (STEMI) paradigm, faces scrutiny regarding its foundational assumptions. While advocated for its simplicity, the paradigm's reliance on specific millimeter-based ST-segment elevation criteria may not consistently reflect the underlying pathophysiologic reality of occlusion myocardial infarction (OMI). This raises questions about the universality of current diagnostic approaches and the interpretation of clinical trial outcomes.

KEY TAKEAWAYS

- **The Pivot:** The STEMI paradigm's fixed ST-elevation criteria are challenged by inconsistencies in diagnosing acute coronary occlusion, suggesting a disconnect from true OMI pathophysiology.
- **The Data:** The STEMI paradigm often fails to reflect the underlying pathophysiologic reality captured in the OMI paradigm.¹
- **The Action:** Clinicians should be aware that current STEMI criteria may not always align with actual coronary occlusion, prompting consideration of broader diagnostic perspectives.

The ST-elevation myocardial infarction (STEMI) paradigm operates on the assumption that specific millimeter-based ST-segment elevation criteria serve as a reliable surrogate for acute coronary occlusion. Proponents of this paradigm advocate for its simplicity and uniformity. However, research indicates that this paradigm often fails to reflect the underlying pathophysiologic reality captured in the Occlusion Myocardial Infarction (OMI) paradigm.¹ This diagnostic inconsistency highlights a potential limitation in the current approach to identifying acute coronary events.¹

Challenges to Established Paradigms

Further examination reveals that the STEMI paradigm's reliance on these specific criteria does not always align with the actual presence of acute coronary occlusion. The OMI paradigm, in contrast, aims to capture this pathophysiologic reality more accurately.¹ The discrepancy between these two paradigms suggests that the simplicity of the STEMI criteria may come at the cost of diagnostic precision.¹

Beyond diagnostic criteria, the interpretation of clinical trial data also warrants re-evaluation. For instance, divergent beta-blocker trials illustrate that randomization alone may be insufficient to establish universal causality.³ This implies that even well-designed randomized controlled trials may produce outcomes that are not universally applicable, challenging the illusion of universal causality in evidence-based medicine.³ The variability in trial results, despite rigorous methodology, underscores the complexity of applying findings broadly across diverse patient populations and clinical contexts.³

The increasing integration of artificial intelligence (AI) in medicine also presents a complex risk/benefit profile, particularly in fields such as toxicology.² While AI offers potential advancements, its application requires careful analysis to avoid what has been termed 'AI snake oil.'² This cautionary perspective suggests that the enthusiasm for new technologies must be tempered with rigorous evaluation of their true efficacy and potential risks, ensuring that they genuinely improve patient outcomes rather than merely adding layers of complexity or unverified claims.² The critical assessment of AI tools, similar to the re-evaluation of diagnostic paradigms and trial interpretations, is essential for maintaining the integrity of evidence-based practice.²

CLINICAL IMPLICATIONS

The continued reliance on the STEMI paradigm, despite its documented inconsistencies with actual coronary occlusion, presents a significant challenge for emergency physicians and cardiologists. When millimeter-based ST-elevation criteria do not accurately reflect the underlying OMI, patient management decisions, particularly regarding reperfusion strategies, may be suboptimal. This calls for a re-evaluation of guideline bodies' recommendations, urging them to consider the OMI paradigm's emphasis on pathophysiologic reality over simplified electrocardiographic thresholds. The current approach risks both over-treatment in some cases and delayed intervention in others, directly impacting patient morbidity and mortality.

The broader implication extends to the pharmaceutical industry and its development of therapies. If clinical trials, even randomized ones, cannot establish universal causality, as seen with beta-blocker trials, then the generalizability of drug efficacy becomes questionable. This necessitates a more nuanced interpretation of trial results, moving beyond headline statistics to understand the specific patient populations and contexts in which a treatment truly offers benefit. Companies developing new cardiovascular drugs, for example, must be prepared to demonstrate efficacy across a wider range of presentations, rather than relying solely on criteria that may not capture the full spectrum of disease.

For patients, the illusion of evidence-based medicine means that what appears to be a clear-cut diagnosis or universally effective treatment may, in reality, be far more complex. The promise of AI in toxicology, for instance, must be approached with caution; the risk of 'AI snake oil' means that patients could be exposed to unverified or even harmful diagnostic or therapeutic recommendations. Clinicians have a responsibility to communicate these nuances transparently, managing patient expectations and advocating for diagnostic and therapeutic approaches that are truly evidence-based, rather than merely evidence-appearing.

EDITORIAL & AI STANDARDS

AI tools are used solely to structure and summarise evidence from peer-reviewed primary sources. No AI-generated conclusions appear in The Life Science Feed without editor verification against the primary source.

REFERENCES

1. Frick WH, Smith SW. The illusion of simplicity: Diagnostic inconsistencies within the STEMI paradigm. J Electrocardiol. 2026;42287922. doi:10.1016/j.jelectrocard.2026.154392

2. Hartung T, Rao M, Behl M. AI snake oil? A risk/benefit analysis for toxicology. *Front Artif Intell.* 2026;42238196. doi:10.3389/frai.2026.1779276
 3. Omerovic E, Råmunddal T, Jha S. When Randomization Is Not Enough: Divergent Beta-Blocker Trials and the Illusion of Universal Causality. *Eur Heart J Cardiovasc Pharmacother.* 2026;42219422. doi:10.1093/ehjcvp/pvag037
-

LICENCE & RIGHTS

© 2026 The Life Science Feed. All rights reserved. This content is produced for medical education purposes. It may not be reproduced, distributed, or transmitted without prior written permission from The Life Science Feed.

MEDICAL DISCLAIMER

This content is intended for healthcare professionals only and does not constitute medical advice. Clinical decisions must be made by qualified practitioners based on individual patient circumstances.

FOLLOW THE LIFE SCIENCE FEED

Instagram @lifesciencefeed **LinkedIn** linkedin.com/company/thelifesciencefeed

thelifesciencefeed.com