

Fibromyalgia: Is the pain real if the tissue damage isn't?

Written by The Life Science Feed Editorial Team · Published 1 October 2026 · Edited by Mara Voss

Fibromyalgia presents a persistent clinical conundrum: patients report widespread, debilitating pain, yet objective signs of peripheral tissue damage are consistently absent. This disconnect has historically complicated diagnosis and management, often leading to skepticism about the condition's biological basis. But, emerging evidence points towards a re-evaluation, positioning fibromyalgia as a disorder primarily rooted in central nervous system dysfunction.

KEY TAKEAWAYS

- **The Pivot:** Fibromyalgia is now largely understood as a central nervous system disorder, not a peripheral tissue pathology.
- **The Data:** Neuroimaging and neurochemical studies consistently show altered pain processing and neurotransmitter imbalances in the brain.
- **The Action:** Clinicians should focus on therapies targeting central sensitisation and neuropathic pain mechanisms, moving beyond anti-inflammatory or musculoskeletal approaches.

Fibromyalgia is characterised by chronic widespread pain, fatigue, sleep disturbances, and cognitive dysfunction. The hallmark of the condition is diffuse musculoskeletal pain that cannot be attributed to inflammation or structural damage in the affected tissues. This absence of peripheral pathology has long been a source of frustration for both patients and clinicians, as standard diagnostic tools like X-rays, MRI, and blood tests for inflammation typically yield normal results. The diagnostic criteria, therefore, rely heavily on subjective reporting of pain and associated symptoms, making it a diagnosis of exclusion.

The patient population affected by fibromyalgia is predominantly female, with symptoms often developing after a physical trauma, infection, or significant psychological stress, though it can also emerge without a clear trigger. The pain is often described as a constant dull ache, but can also manifest as burning, shooting, or throbbing sensations. The widespread nature of the pain, affecting both sides of the body and above and below the waist, is a key diagnostic feature. The absence of clear tissue damage differentiates it from inflammatory arthropathies or localised neuropathies.

The Central Sensitisation Hypothesis

The prevailing hypothesis for fibromyalgia's pathophysiology centers on central sensitisation, a phenomenon where the central nervous system becomes hypersensitive to pain signals. This involves an amplification of pain processing within the brain and spinal cord, leading to an exaggerated response to stimuli that would normally not be painful (allodynia) or an increased response to painful stimuli (hyperalgesia). This mechanism explains why patients experience widespread

pain without peripheral tissue damage; the problem lies in how the brain interprets and processes sensory input, not in the peripheral tissues themselves.

Neuroimaging studies, particularly functional magnetic resonance imaging (fMRI), have provided compelling evidence for altered pain processing in fibromyalgia patients. These studies consistently show abnormal activation patterns in brain regions involved in pain perception, emotion, and cognition. For instance, areas like the insula, anterior cingulate cortex, and prefrontal cortex, which form part of the 'pain matrix', exhibit heightened activity in response to noxious stimuli. This hyperactivity suggests a dysregulation in descending pain inhibitory pathways, which normally modulate and dampen pain signals.

Beyond imaging, neurochemical investigations have identified imbalances in neurotransmitter systems implicated in pain modulation. Patients with fibromyalgia often exhibit lower levels of inhibitory neurotransmitters, such as gamma-aminobutyric acid (GABA), and elevated levels of excitatory neurotransmitters, like substance P and glutamate, in the cerebrospinal fluid. These neurochemical alterations contribute to the hyperexcitability of pain pathways. The role of these imbalances in the broader symptom complex, including fatigue and cognitive issues, is also under active investigation, suggesting a more global central nervous system dysregulation.

The understanding of central sensitisation has shifted the therapeutic focus from peripheral pain relief to modulating central nervous system activity. This is why traditional analgesics, such as non-steroidal anti-inflammatory drugs (NSAIDs), often provide limited benefit in fibromyalgia, as they primarily target peripheral inflammation. Instead, treatments that influence neurotransmitter systems or neural pathways involved in pain processing have shown more promise. This includes certain antidepressants and anticonvulsants, which act on serotonin, norepinephrine, and GABA pathways.

Neuroendocrine and Immune System Dysregulation

Fibromyalgia is not solely a pain disorder; it is a complex syndrome with systemic manifestations, including profound fatigue, sleep disturbances, and cognitive difficulties often described as 'fibro fog'. These symptoms point to broader dysregulation within the central nervous system, extending beyond pain processing pathways. The relationship between the nervous, endocrine, and immune systems appears to be a critical factor in the disease's pathogenesis, further supporting its classification as a central nervous system disorder.

Dysfunction of the hypothalamic-pituitary-adrenal (HPA) axis, the body's central stress response system, is frequently observed in fibromyalgia patients. Studies have reported altered cortisol levels, blunted cortisol responses to stress, and abnormal diurnal cortisol rhythms. This HPA axis dysregulation can contribute to chronic fatigue, sleep disturbances, and heightened stress sensitivity, all common features of the condition. The HPA axis is intricately linked to both pain perception and immune function, creating a complex feedback loop that can perpetuate symptoms.

The immune system also shows signs of dysregulation in fibromyalgia, though not in the form of overt inflammation. Instead, there is evidence of altered cytokine profiles and immune cell activation, suggesting a state of chronic low-grade neuroinflammation or immune system imbalance. For example, elevated levels of pro-inflammatory cytokines such as IL-6 and IL-8 have been detected in some patients, alongside reduced anti-inflammatory cytokines. This subtle immune activation within the central nervous system could contribute to pain sensitisation and other neurological symptoms. The question of immune system modulation is a broader one across many conditions.

Sleep disturbances are almost universal in fibromyalgia, with patients often experiencing non-restorative sleep and

alpha-wave intrusion into non-REM sleep. This abnormal sleep architecture can exacerbate pain, fatigue, and cognitive dysfunction, creating a vicious cycle. The brain's ability to process and consolidate memories, as well as clear metabolic waste products, is impaired during disrupted sleep. This further reinforces the idea that central nervous system dysfunction is at the core of the syndrome, affecting multiple interconnected physiological processes. For more on this, consider our previous coverage on cyclobenzaprine and sleep improvement in fibromyalgia.

The cognitive symptoms, including difficulties with memory, concentration, and information processing, are particularly distressing for patients. These 'fibro fog' symptoms are consistent with the observed neurochemical and functional brain changes, particularly in areas responsible for executive function and attention. The cumulative effect of chronic pain, fatigue, and sleep deprivation likely contributes to these cognitive deficits, highlighting the systemic impact of central nervous system dysregulation. Understanding these interconnected systems is essential for developing comprehensive management strategies that improve patient outcomes.

Therapeutic Implications and Future Directions

The understanding of fibromyalgia as a central nervous system disorder has profound implications for its management. Current guideline recommendations for fibromyalgia often include a multimodal approach, combining pharmacological and non-pharmacological therapies. Pharmacological interventions primarily target neurotransmitter systems involved in pain and mood regulation. These include serotonin-norepinephrine reuptake inhibitors (SNRIs) like duloxetine and milnacipran, which increase levels of serotonin and norepinephrine in the brain, thereby enhancing descending pain inhibition. Pregabalin and gabapentin, alpha-2-delta ligands, are also used to modulate calcium channels and reduce excitatory neurotransmitter release, dampening neuronal hyperexcitability.

Non-pharmacological treatments are equally critical and often form the cornerstone of long-term management. Cognitive behavioral therapy (CBT) helps patients reframe their perception of pain and develop coping strategies, addressing the psychological components of chronic pain. Graded exercise therapy, tailored to individual tolerance, can improve physical function, reduce pain, and alleviate fatigue, despite initial apprehension from patients. Other approaches like mindfulness, acupuncture, and hydrotherapy are also explored, aiming to modulate central pain processing and improve overall well-being. A comprehensive reference like the Oxford Handbook of Clinical Medicine can be invaluable for navigating these complex, multi-system conditions.

Despite these advances, significant unmet needs remain. Many patients do not achieve adequate symptom control with existing therapies, and treatment responses can be highly variable. The heterogeneity of fibromyalgia symptoms suggests that different underlying mechanisms may be at play in different individuals, necessitating a more personalised medicine approach. Future research is focusing on identifying specific biomarkers that could predict treatment response or stratify patients into distinct subgroups based on their underlying pathophysiology. This could lead to more targeted and effective interventions.

The development of novel therapeutic targets is also underway, exploring pathways beyond the traditional neurotransmitter systems. For instance, research into glial cell activation, neuroinflammation, and mitochondrial dysfunction is gaining traction. These areas represent potential new avenues for intervention, aiming to address the root causes of central sensitisation and broader central nervous system dysregulation. The ongoing challenge is to translate these mechanistic insights into clinically meaningful treatments that can improve the quality of life for patients living with this debilitating condition. The field needs to move beyond symptomatic relief to address the underlying neurobiological

changes, a complex task given the diffuse nature of the disorder.

The absence of peripheral tissue damage does not mean the pain is not real. It means we have been looking in the wrong place. Sarah Gellar, Clinical Trials Editor

CLINICAL IMPLICATIONS

The shift in understanding fibromyalgia from a peripheral tissue disorder to a central nervous system disorder is not merely academic; it fundamentally changes how clinicians should approach diagnosis and treatment. GPs and specialists alike must internalise that the absence of objective peripheral findings does not invalidate the patient's experience of pain. Dismissing symptoms because a joint X-ray is clear or inflammatory markers are normal is a disservice to patients and a misunderstanding of the current evidence.

This re-framing means that therapies targeting inflammation or structural damage are largely ineffective. Instead, the focus must be on modulating central pain processing. This includes judicious use of agents like SNRIs and alpha-2-delta ligands, alongside robust non-pharmacological interventions such as CBT and graded exercise. The challenge lies in integrating these complex, often time-intensive, approaches into routine clinical practice, especially given the resource constraints in many healthcare systems.

For patients, this evolving understanding offers both validation and a clearer path forward. Knowing that their pain has a neurobiological basis, even if not visible on a scan, can reduce the psychological burden of being disbelieved. It also empowers them to engage with therapies that target the central nervous system, rather than endlessly pursuing treatments for non-existent peripheral pathology. The industry, in turn, needs to invest in developing novel compounds that specifically address the identified neurochemical imbalances and central sensitisation mechanisms, moving beyond repurposed antidepressants.

EDITORIAL & AI STANDARDS

AI tools are used solely to structure and summarise evidence from peer-reviewed primary sources. No AI-generated conclusions appear in The Life Science Feed without editor verification against the primary source.

REFERENCES

1. Kortright-Maldonado K, Deanda-Quintero J, Martínez-Lavin M. Fibromyalgia: a real phantom pain? Clin Rheumatol. 2026;45(9):5941-5944. doi:10.1007/s10067-026-08331-4
2. Vittersø AD, Halicka M, Buckingham G, Proulx MJ, Bultitude JH. The sensorimotor theory of pathological pain revisited. Neurosci Biobehav Rev. 2022;139:104735. doi:10.1016/j.neubiorev.2022.104735
3. Banfi G, Lombardi G, Colombini A, Melegati G. Whole-body cryotherapy in athletes. Sports Med. 2010;40(6):509-17. doi:10.2165/11531940-000000000-00000
4. Guymer EK, Littlejohn GO. Fibromyalgia. Diagnosis and management. Australas Chiropr Osteopathy. 2002;10(2):81-4. PMID:17987178

LICENCE & RIGHTS

© 2026 The Life Science Feed. All rights reserved. This content is produced for medical education purposes. It may not be reproduced, distributed, or transmitted without prior written permission from The Life Science Feed.

MEDICAL DISCLAIMER

This content is intended for healthcare professionals only and does not constitute medical advice. Clinical decisions must be made by qualified practitioners based on individual patient circumstances.

FOLLOW THE LIFE SCIENCE FEED

Instagram @lifesciencefeed **LinkedIn** linkedin.com/company/thelifesciencefeed

thelifesciencefeed.com