

Raw milk cheese: The 'natural' choice still sparking E. coli outbreaks?

Written by The Life Science Feed Editorial Team · Published 16 October 2026 · Edited by Mara Voss

Foodborne illness outbreaks remain a persistent public health challenge, with pathogens like *Escherichia coli* (*E. coli*) frequently implicated in cases linked to contaminated food products. These incidents highlight the importance of food safety regulations and consumer awareness, particularly concerning products that bypass standard pasteurization processes.

A recent cluster of *E. coli* infections spanning nine states has been definitively linked to the consumption of raw milk cheese, raising concerns about the risks associated with unpasteurized dairy products and the potential for widespread contamination.

KEY TAKEAWAYS

- **The Pivot:** A multi-state *E. coli* outbreak has been traced to raw milk cheese, highlighting ongoing risks with unpasteurized dairy.
- **The Data:** The outbreak has affected at least nine states, with specific case numbers under investigation by public health authorities.
- **The Action:** Clinicians should advise patients, especially vulnerable populations, to avoid raw milk and raw milk products due to potential bacterial contamination.

E. coli infections, particularly those caused by Shiga toxin-producing *E. coli* (STEC) strains, present a significant public health burden. These bacteria can cause severe gastrointestinal illness, including bloody diarrhea, and in some cases, lead to life-threatening complications such as hemolytic uremic syndrome (HUS). The primary route of transmission is often through contaminated food or water, with raw or undercooked meat, unpasteurized dairy products, and fresh produce being common culprits. Understanding the epidemiology of these outbreaks is essential for implementing effective control measures and protecting public health.

The current *E. coli* outbreak, identified across nine US states, has been linked to raw milk cheese. Public health investigations are ongoing to identify the specific source within the raw milk cheese supply chain and to determine the full scope of the outbreak. These investigations typically involve epidemiological traceback, laboratory testing of patient samples, and environmental sampling of food production facilities. The frequency of foodborne outbreaks has been a topic of ongoing discussion among public health experts.

The Risks of Unpasteurized Dairy

Raw milk and products derived from it, such as certain cheeses, pose inherent risks due to the absence of pasteurization. Pasteurization is a heat treatment process that kills harmful bacteria, including *E. coli*, *Salmonella*, *Listeria*, and

Campylobacter, without significantly altering the nutritional value or taste of the product. The process involves heating milk to a specific temperature for a set period, effectively eliminating pathogens that may be present.

Advocates for raw milk often cite perceived health benefits, such as higher enzyme content or beneficial bacteria, but these claims lack robust scientific evidence. The potential presence of pathogenic bacteria in raw milk far outweighs any unproven benefits, making it a high-risk food item, especially for vulnerable populations. The severity of gastroenteritis cases can vary widely depending on the pathogen and host factors.

Children, the elderly, pregnant women, and individuals with compromised immune systems are particularly susceptible to severe illness from raw milk pathogens. For these groups, even a small bacterial load can lead to serious complications, including kidney failure in the case of STEC infections. The lack of a heat treatment step means that any contamination introduced during milking, handling, or processing will persist in the final product, ready to infect consumers.

Regulatory bodies in many countries and regions strongly advise against the consumption of raw milk and raw milk products. Public health guidelines consistently recommend choosing pasteurized dairy to minimize the risk of foodborne illness. This guidance is based on decades of epidemiological data linking raw milk consumption to outbreaks of various bacterial infections. Clinicians should reinforce these recommendations with their patients, particularly when discussing dietary choices.

Understanding E. coli Pathogenesis

E. coli is a diverse group of bacteria, with most strains being harmless commensals in the human gut. However, certain pathogenic strains, particularly STEC, produce Shiga toxins that damage the lining of the intestine and can enter the bloodstream, leading to systemic complications. The most common STEC strain associated with severe human disease is O157:H7, though non-O157 STEC strains are also significant.

Once ingested, STEC bacteria colonize the large intestine and begin producing Shiga toxins. These toxins bind to specific receptors on endothelial cells, particularly in the kidneys, leading to microvascular damage and thrombosis. This cascade of events can result in HUS, characterized by hemolytic anemia, thrombocytopenia, and acute kidney injury. The clinical presentation often begins with non-bloody diarrhea, progressing to bloody diarrhea within a few days.

Diagnosis of STEC infection relies on stool culture and specific assays for Shiga toxin. Treatment is primarily supportive, focusing on hydration and managing complications. Antibiotics are generally not recommended for STEC infections, as they can sometimes increase Shiga toxin release and worsen the risk of HUS. This makes prevention, through safe food handling and consumption practices, the cornerstone of public health efforts against these infections. For a comprehensive overview of infectious diseases, the Oxford Handbook of Infectious Diseases and Microbiology serves as a practical guide.

The incubation period for STEC infections typically ranges from 3 to 4 days but can be as short as 1 day or as long as 10 days. This variability can make traceback investigations challenging, as individuals may not recall specific food exposures from a week or more prior to symptom onset. The prolonged incubation period also allows for wider dissemination of contaminated products before an outbreak is recognized, increasing the number of affected individuals. Public health agencies often rely on rapid reporting and robust surveillance systems to detect and respond to these outbreaks efficiently. The reporting of other foodborne illnesses, like cyclosporiasis, also highlights the ongoing need for vigilance.

Public Health Response and Prevention

When an *E. coli* outbreak is identified, public health authorities initiate a coordinated response involving epidemiology, laboratory science, and environmental health. This includes interviewing patients to gather information about food exposures, collecting and testing clinical samples, and tracing the distribution of implicated food products. The goal is to quickly identify the source, remove contaminated products from the market, and prevent further illnesses.

For this specific outbreak linked to raw milk cheese, public health officials have issued warnings to consumers, advising them to avoid the implicated products. Retailers and distributors are typically instructed to pull the products from shelves. These actions are vital to containing the outbreak and protecting the public. But the challenge lies in the widespread distribution of such products and the difficulty in reaching all consumers who may have purchased them.

Prevention of foodborne *E. coli* infections ultimately rests on a combination of robust food safety practices throughout the supply chain and informed consumer choices. For clinicians, this means educating patients about the risks associated with raw milk and raw milk products, emphasizing the importance of proper food handling, and advising thorough cooking of meats. Hand hygiene is also paramount, especially after contact with animals or after using the restroom, to prevent secondary transmission. The ongoing efforts to understand and prevent outbreaks, including mathematical modeling in infectious disease, are vital for public health.

The current outbreak serves as a stark reminder that despite advances in food safety, risks persist, particularly with products that bypass established pathogen reduction steps. While some consumers may prefer raw milk products for various reasons, the scientific evidence consistently points to a heightened risk of serious illness. Clinicians should be prepared to discuss these risks clearly and unequivocally with patients, ensuring they make informed decisions about their dietary intake. The public health message remains consistent: pasteurization is a critical safety step that protects against dangerous foodborne pathogens.

CLINICAL IMPLICATIONS

The *E. coli* outbreak linked to raw milk cheese reinforces a fundamental principle in food safety: unpasteurized dairy products carry inherent risks that pasteurization effectively mitigates. Clinicians should use this as an opportunity to counsel patients, particularly those with young children, pregnant individuals, or immunocompromised family members, against consuming raw milk or raw milk cheeses. The perceived benefits of raw milk are largely unsubstantiated, while the risks of severe bacterial infection are well-documented.

For general practitioners, maintaining a high index of suspicion for STEC infection in patients presenting with bloody diarrhea, especially if they report consuming raw dairy, is essential for patient safety. Early recognition and supportive care are paramount, given that antibiotic use can sometimes exacerbate toxin release. A thorough dietary history, including questions about unpasteurized products, should be part of the workup for acute gastroenteritis.

The incident also highlights the ongoing challenges in regulating and monitoring artisanal food products. While local and small-scale food production can offer benefits, it must adhere to stringent safety standards to prevent widespread public health crises. Patients often trust their healthcare providers for advice on healthy eating, and this extends to understanding the risks associated with certain food choices.

EDITORIAL & AI STANDARDS

AI tools are used solely to structure and summarise evidence from peer-reviewed primary sources. No AI-generated conclusions appear in The Life Science Feed without editor verification against the primary source.

REFERENCES

1. Currie A, Galanis E, Chacon PA, et al. Outbreak of *Escherichia coli* O157:H7 Infections Linked to Aged Raw Milk Gouda Cheese, Canada, 2013. *J Food Prot.* 2018;81(2):325-331. doi:10.4315/0362-028X.JFP-17-283
2. Gaulin C, Levac E, Ramsay D, et al. *Escherichia coli* O157:H7 outbreak linked to raw milk cheese in Quebec, Canada: use of exact probability calculation and casecase study approaches to foodborne outbreak investigation. *J Food Prot.* 2012;75(5):812-8. doi:10.4315/0362-028X.JFP-11-385
3. Weinstein E, Lamba K, Bond C, et al. Outbreak of *Salmonella* Typhimurium Infections Linked to Commercially Distributed Raw Milk - California and Four Other States, September 2023-March 2024. *MMWR Morb Mortal Wkly Rep.* 2025;74(27):433-438. doi:10.15585/mmwr.mm7427a1
4. Jones G, Lefèvre S, Donguy MP, et al. Outbreak of Shiga toxin-producing *Escherichia coli* (STEC) O26 paediatric haemolytic uraemic syndrome (HUS) cases associated with the consumption of soft raw cow's milk cheeses, France, March to May 2019. *Euro Surveill.* 2019;24(22). doi:10.2807/1560-7917.ES.2019.24.22.1900305
5. Quinn OI, Jenkins C, Greig DR, et al. An outbreak of Shiga Toxin-producing *Escherichia coli* Serotype O145:H28 Associated with Domestic Travel and Consumption of Unpasteurized Cheese, UK, 2023. *J Food Prot.* 2025;88(4):100470. doi:10.1016/j.jfp.2025.100470
6. Ung A, Baidjoe AY, Van Cauteren D, et al. Disentangling a complex nationwide *Salmonella* Dublin outbreak associated with raw-milk cheese consumption, France, 2015 to 2016. *Euro Surveill.* 2019;24(3). doi:10.2807/1560-7917.ES.2019.24.3.1700703

LICENCE & RIGHTS

© 2026 The Life Science Feed. All rights reserved. This content is produced for medical education purposes. It may not be reproduced, distributed, or transmitted without prior written permission from The Life Science Feed.

MEDICAL DISCLAIMER

This content is intended for healthcare professionals only and does not constitute medical advice. Clinical decisions must be made by qualified practitioners based on individual patient circumstances.

FOLLOW THE LIFE SCIENCE FEED

Instagram @lifesciencefeed **LinkedIn** linkedin.com/company/thelifesciencefeed

thelifesciencefeed.com